

*Research Article***Comprehensive Analysis of Complete Blood Count in ICU Patients at a Tertiary Medical Facility*****Shadma Siddiqui, ** Pinki Vishwakarma**

* School of Paramedical Sciences, SAM Global University, Raisen, M.P.

** Department of Paramedical Sciences, NIMS University, Jaipur, RJ

Article information**ABSTRACT****Volume: 1****Issue: 1****Page No: 56-62****Received: 17.07.2024****Accepted: 22.7.2024****Published: 26.09.2024****DOI No.:****Corresponding Author:**

Dr. Shadma Siddiqui
Professor, SAM Global
University, Raisen (M.P),
Email:
shadma.10aug@gmail.com
Mobile:- 9993811145

Keywords:

Complete Blood Count,
ICU, Haematological
Analysis, Critical Care,
Blood Parameters, Patient
Monitoring, Diagnostic
Biomarkers, Prognostic
Indicators.

The complete blood count (CBC) is a critical diagnostic tool in intensive care settings, providing valuable insights into haematological abnormalities that may influence patient prognosis. This study examines CBC variations in ICU patients at a tertiary healthcare centre, focusing on key parameters such as white blood cell (WBC) count, red blood cell (RBC) levels, haemoglobin concentration, and platelet count. Elevated WBC levels were often indicative of systemic infections, while anaemia and thrombocytopenia were prevalent in critically ill patients, correlating with increased mortality risks. Gender-based differences were observed, with male patients exhibiting higher haemoglobin levels than females. A strong association between severe CBC abnormalities and higher ICU mortality rates was identified, underscoring the importance of timely intervention. Statistical analyses confirmed significant deviations in CBC values from healthy reference ranges, reinforcing the role of routine haematological monitoring in critical care. Findings from this study highlight the need for integrated diagnostic approaches and tailored therapeutic strategies to enhance patient survival. Future research should explore longitudinal CBC trends and their predictive value in intensive care outcomes. This study emphasizes the necessity for continuous monitoring and personalized treatment to optimize ICU patient care.

INTRODUCTION

Intensive Care Units (ICUs) serve as specialized hospital departments where patients with life-threatening conditions receive constant medical supervision and treatment. Critical care medicine relies on a variety of diagnostic tools, among which the complete blood count (CBC) is one of the most frequently utilized tests. CBC analysis provides crucial insights into a patient's overall health status by evaluating key blood components such as white blood cells (WBC), red blood cells (RBC), and platelets.

This study examines CBC results from ICU patients, assessing their variations and significance in disease progression and management. A detailed statistical evaluation has been conducted, considering gender-based differences, normal distribution assessments, and significant deviations from standard values.

METHODOLOGY

A cohort of 100 healthy volunteers (56 males and 44 females) served as the control group for establishing normal CBC ranges. The study population included critically ill patients aged between 22 and 72 years who were admitted to the ICU. Statistical analysis was performed using the Shapiro-Wilk test to determine data distribution. For normally distributed parameters, results were presented as mean \pm standard deviation (SD), whereas non-normally distributed variables were reported

using median and interquartile range (IQR). Gender-based differences were analyzed through the Mann-Whitney U test, with statistical significance set at $p < 0.05$.

Exclusion criteria included patients undergoing cardiac surgery, obstetric cases, and studies only available in abstract form or in non-English publications. Citations were screened independently by two reviewers to ensure accuracy and relevance.

RESULTS AND DISCUSSION

This study identified significant variations in CBC values among ICU patients, reflecting the impact of critical illness on haematological parameters. Findings include:

White Blood Cells (WBC) Variations

Elevated WBC counts were observed in patients with systemic infections and inflammatory responses, supporting their role as biomarkers for sepsis and immune dysfunction. Leucocytosis, commonly present in ICU patients, suggests an active immune response, whereas leukopenia indicates immunosuppression or severe systemic involvement. These variations play a crucial role in determining treatment protocols, including antibiotic administration and infection control measures.


Red Blood Cells (RBC) & Haemoglobin Trends

Anaemia was prevalent among critically ill patients, often linked to underlying chronic conditions, nutritional deficiencies, or blood

loss due to medical interventions. A low RBC count and decreased haemoglobin levels were associated with poor oxygen-carrying capacity, leading to increased risks of organ dysfunction. Blood transfusions, iron supplementation, and erythropoiesis-stimulating agents were frequently utilized interventions for managing anaemia in ICU settings.

Platelet Count and Coagulation Concerns


Thrombocytopenia was a common finding, correlating with an increased risk of bleeding complications and poor clinical outcomes. Platelets play a pivotal role in haemostasis and immune function, and their depletion in critically ill patients could result from sepsis, disseminated intravascular coagulation (DIC), or drug-induced effects. Monitoring platelet levels is crucial for managing bleeding risks and implementing appropriate therapeutic strategies, such as platelet transfusions and coagulation factor replacement therapies.



Graph-I: compares WBC, RBC, and platelet counts between healthy individuals and ICU patients. ICU patients show significant deviations, particularly in WBC (higher due to infections) and platelets (lower due to critical conditions).

Comparative Gender-Based Analysis

Gender differences were analyzed to determine CBC variations among male and female ICU patients. Statistical tests revealed significant disparities in haemoglobin levels, with male patients exhibiting slightly higher values than females. However, WBC and platelet counts showed minimal variation between genders, reinforcing the necessity of individualized patient assessments rather than relying solely on generalized reference ranges.




Graph-II: illustrates gender-based differences in CBC parameters among ICU patients. Male patients have slightly higher haemoglobin levels, while WBC and platelet counts show minor variations between genders.

Correlation Between CBC and ICU Mortality

This study also investigated the predictive value of CBC variations in determining ICU mortality rates. Patients with extreme deviations in WBC, RBC, and platelet counts were found to have higher mortality risks. Early recognition of these abnormalities and

timely medical interventions were instrumental in reducing ICU mortality and improving



patient survival rates.

Graph-III: shows the correlation between CBC abnormality severity and ICU mortality rates. As abnormalities become more severe, mortality rates increase significantly, emphasizing the importance of early detection and intervention.

ETHICAL CONSIDERATIONS AND RESEARCH LIMITATIONS

Ethical approval was obtained from relevant institutional review boards, and informed consent was secured from all participants. This study acknowledges potential limitations, including sample size constraints and the exclusion of specific patient populations that could influence findings. Future research should focus on expanding the dataset and incorporating additional biomarkers to enhance diagnostic accuracy.

CONCLUSION

This research underscores the critical role of CBC in managing ICU patients, highlighting its significance in detecting abnormalities and guiding treatment decisions. The study

advocates for integrating CBC with advanced haematological markers and continuous monitoring to optimize patient care and improve survival outcomes. Further investigations are recommended to explore the predictive value of CBC variations in long-term ICU patient prognosis.

FUTURE RESEARCH DIRECTIONS

Given the findings of this study, further research should explore:

Longitudinal Analysis: Tracking CBC trends over extended ICU stays to evaluate progressive changes and recovery indicators.

Multi-Parameter Integration: Combining CBC data with biomarkers like C-reactive protein (CRP) and procalcitonin for enhanced diagnostic accuracy.

Machine Learning Applications: Utilizing artificial intelligence to predict ICU patient outcomes based on haematological patterns.

ACKNOWLEDGMENTS

The authors express their gratitude to colleagues, faculty members, and supporting staff for their contributions to this study. Special appreciation is extended to the research participants for their valuable cooperation.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest related to this research.

REFERENCES

1. Abel N Kho, MD MS: siuHui PHD
Joe G Kesterson, BS: Clement J
MD journal of hospital medicine
2007: 2nd edition (1) 6-12.
2. American Academy of
Dermatology (AAD). (2020)
"Cosmetics and Skin Care:
Allergies and Irritations." Retrieved
from [AAD Website]
(<https://www.aad.org>)
3. Banck M, Walther S,
KarlstromG,Noblin T,
F,SamuelssonC. There are more
male than female patients within
intensive care. But it is still unclear
whether there are gender
inequasilities in Swedish intensive
care or not. Lakartiningen.
2014;111 (9-10):388-90
4. Barr, Fraser GL, Puntillo K, et
al.:Anurican college of critical care
medicine:Clinical guidelines for the
management of pain agitation and
deliriunl in adult patients in the
ICU. Critical care medicine
2013;41; 263-306.
5. Brophy DF et al.Critical illness in
ICU patients Anemia 2012
6. Cong YL, jin DM, Wang HL,
Okada T, Peng ZH. Establishing the
reference range of venous blood
measured by automated
haematology analyzer in Chinese
adults. Zhonghua Yi XueZaZhi.
2003; 83:1201-1205.
7. Deutschman, Clifford S: Neligan,
Patrick j. (2010). Evidence- based
practice of Critical Care Elsevier
Health Sciences: ISBN
1416054766. Retrived 2015-04-30.
8. Diagnostic tools and therapeutic
end-points in critical care medicine:
Limitations and concepts Journal of
Nursing Education and Practice,
2014, Vol. 4, No. 7
9. Effect of Patient Sex on Intensive
Care Unit Survival, Hugo Romo,
MD; Andre Kajdacsy-
BallaAmaral, MD; Jean-Louis
Vincent, MD, PHD, Arch Intern
Med. 2004; 164(1): 61-65 doi:10.
1001/ archinte. 164.61
10. ElieAzoulay, Giuseppe citerio, Jan
Bakker, Gleen Hernandez, Matteo
Bassetti and 10 more,
(10.1007/S00134-014-3217-8)
March 2014, volume 40,issue 3, pp
305-319.
11. Evaluation of patients with
Leukocytosis, LYRAD K. RILEY,
MD, and JEDDA RUPERT, MD,
Am Fam physician. 2015 December
1;92 (11): 1004-1011.
12. Fischbach FT, Dunning MB 3rd
edition (2009). Manual of

- laboratory and Diagnostic Test, 8th edition Philadelphia: Lippincott Williams and Wilkins.
13. Guidelines to practice of EMERGENCY MEDICINE, Narayan H Nayak, M.S. (General surgery), M.D. FACEP, second edition 2008 Elsevier 77-
 14. Hatden SJ et al: Am J Respir Crit Care Med. 2012.
 15. International Journal of Pharma and Bio Sciences Research Article DR. RAJENDRA GUPTA Int J Pharm Bio Sci 2015 July; 6(3): (P) 325 – 329.
 16. Lutan, Vasile. Fiziopatologic medicală. Vol. 2, 31.3.2.1 Leucocitoză <http://library.usmf.md/ebooks?Key>
 17. Pediatrie polska (optimizing use of CBC) Bruce M. Camitta, Rebecca Jean Slye (jan-feb 2012, volume 8 (1): 72-77, doi:10
 18. Porth, Carol Mattson (2011), “White blood cell response”, Essentials of Pathophysiology: Concepts of Altered Health States (3rd ed.), Philadelphia: Wolters Kluwer Health/ Lippincott Williams & Wilkins, pp 64-65, ISBN 978-1-58255-724-3, retrieved 13 November 2011
 19. Prevalence of unnecessary laboratory tests and related avoidable costs in intensive care unit, Anderson Magalhães Oliveira¹; Marcio Vasconcelos Oliveira²; Claudio Lima Souza² J Bras Patol Med Lab, volume 50, n. 6, pp 410-416, dezembro 2014.
 20. Principles of internal medicine, DENNIS L. KASPER, MD, ANTHONY S. FAUCI, MD, DAN L. LONGO, MD, EUGENE BRAUNWALD, MD, STEPHEN L. HAUSER, MD, J. LARRY JAMESON, MD, PHD, HARRISON'S volume II, 16th edition, McGraw-Hill medical publishing Division; 1581-1586.
 21. Rogers, Kara, ed. (2011), “Leukocytosis Definition”, “Blood: Physiology and Circulation, Chicago: Britannica Educational Publishing, p. 198, ISBN 978-1-61530-250-5, RETRIVED 12 November 2011
 22. Schwartz, M. William, ed (2013), Leukocytosis, The 5- Minute Pediatrics Consult (3rd ed), Philadelphia: Lippincott Williams & Wilkins, p54, ISBN 0—7817-3539-4 RETRIVED 12 November 2011
 23. Siamak N. Nabili. MD, MPH Meeissa Conrad stoppler, MD, chief

- complete blood count (CBC) [www.Medicine.com/home/complete blood count index](http://www.Medicine.com/home/complete-blood-count-index).
24. Silva JM Jr, Oliveira AM, Segura JL, Ribeiro MH, Sposito Cn, Toledo DO, Rezende E, Malbouisson LM. A large venous-arterial PCO₂ (2) is associated with poor outcomes in surgical patients *Anesthesiol Res Pract*. 2011;2011: 759-792.
 25. The Free Dictionary >Leukocytosis Citing: Gale Encyclopedia of Medicine, 2008 and The American Heritage Medical Dictionary, 2007
 26. Verso, ML (May 1962).” The Evolution of Blood Counting Techniques”.Read at a meeting of the section of the History of Medicine, First Austrilian Medical Congress 8:149-158. Retrieved 9 September 2013.
 27. Walsh TS et al. moderate to severe anemia during ICU stay *BMJ*. 2010
 28. www.UK/Introduction.aspx intensive care unit-NHS choice.
 29. www.ACIHealth.NSW.Gov.AU common blood and other pathology tests in intensive care unit (ICU).
 30. Year in review in Intensive Care Medicine 2013:II. Sedation ,hematological issues in critically ill patients, March 2014, volume 40,issue 3, pp 305-319.
 31. Zore, Joseph J, ed. (2009): Leukocytosis, Schwartz Clinical Handbook of Pediatrics (4th ed): Philadelphia: Wolters Kluwer/ Lippincott Williams & Wilkins, P 559, ISBN 978-0-7817-7013-2, RETRIVED 12 November 2011